

Risk assessment of atmospheric heavy metals exposure in Baotou, a typical industrial city in northern China

Kexin Li · Tao Liang · Lingqing Wang

Received: 11 March 2015 / Accepted: 1 September 2015 / Published online: 8 September 2015
© Springer Science+Business Media Dordrecht 2015

Abstract Industrial activities have led to serious air pollution and the potentially toxic elements in atmospheric particles can cause various health problems to humans. In this study, inhalable particulate matter (PM₁₀) and fine particles (PM_{2.5}) were collected from four typical sites in Baotou, an industrial city in northern China. We investigated both the mass concentrations of particulate matter and the concentrations of heavy metals (Cr, Ni, Pb, Cd, Cu, Mn, Co, and Zn) in the collected samples. We assessed the public health risks associated with atmospheric heavy metal exposure. The results showed that the mass concentrations of PM_{2.5} and PM₁₀ as well as these heavy metal concentrations varied notably influenced by the prevailing wind directions. Among the studied metals, Zn, Mn, Pb, and Cr were the main metal pollutants in both PM₁₀ and PM_{2.5}. The results of the health assessment showed that the eight heavy metals studied pose significant non-carcinogenic risks and Cr, Cd, and Co pose lifetime lung cancer risks to local residents, especially to children.

Keywords PM₁₀ · PM_{2.5} · Heavy metals · Health risk · Baotou

Introduction

Mining activities are notorious for adverse environmental impacts, including pollution, habitat loss, soil erosion, and geological disasters (Salomons 1995; Klukanová and Rapant 1999; Aguilar et al. 2004; Luís et al. 2011). Among these environmental problems caused by mining activities, the production and dispersion of atmospheric particles has become a great concern in recent years (Kaonga and Kgabi 2011; Chen et al. 2013; Serbula et al. 2014). Atmospheric particulates are generated from numerous sources, both natural and anthropogenic, but in areas near mining, mining operations are considered the largest contributor. The processes of mining, such as crushing, grinding, excavating, smelting, and refining, can produce large quantities of particulate matter (PM), containing dangerously high levels of heavy metals (Csavina et al. 2012).

Inhalable particulate matter (PM₁₀), particularly fine particles (PM_{2.5}), have been shown to cause adverse effects on human health, including asthma, lung cancer, and cardiovascular diseases (Pope et al. 2002; Sanchez et al. 2009). Lung and other organ injuries from atmospheric heavy metal exposure have also been well-documented (Espinosa et al. 2001; Cancio et al. 2008; Leili et al. 2008).

K. Li · T. Liang (✉) · L. Wang (✉)
Key Laboratory of Land Surface Pattern and Simulation,
Institute of Geographical Sciences and Natural Resources
Research, Chinese Academy of Sciences, Beijing 100101,
China
e-mail: liangt@igsnr.ac.cn

L. Wang
e-mail: wanglq@igsnr.ac.cn

Baotou city, one of the most industrialized cities of northwest China, has very poor air quality due to large-scale mining activities in the area (Zhen 2012). Toxic elements are mobilized and released into the air as particle matter from anthropogenic activities and pose significant health risks to local residents. However, $PM_{2.5}$ and PM_{10} of Baotou has only been monitored in very recent years. Moreover, little work has addressed the association between the concentrations of aerosol metallic elements and adverse health impacts on the local residents.

The objective of this study was to investigate both the total mass and heavy metal concentrations of $PM_{2.5}$ and PM_{10} as well as how these concentrations varied spatially throughout the city. Sampling was performed in four representative areas of Baotou, including a mining area (MA), an industrial area, a residential area (RA), and the city center. This study also aimed at estimating the non-carcinogenic health risks and lifetime cancer risks of heavy metals exposure for local residents.

Materials and methods


Description of sampling sites

Baotou is located in the central Inner Mongolia Autonomous Region of China. The city is situated on the Tumochuan and Hetao Plateau, and the Yin

Mountains cross the urban area in the central part. The area borders Mongolia to the north, and in the south, the Yellow River runs through the city west to the east for 22 km. Baotou has a cold, semi-arid, continental monsoon climate. The mean annual temperature is 7.2 °C. The annual average rainfall is approximately 310 mm, with the majority mostly occurring in July and August when the southeast monsoon carries rainwater into the province. The prevailing wind direction is northwest, with an average wind speed of 1.2 m s^{-1} . Due to the arid climate and large temperature differences between day and night, dust storms frequently occur in this area, especially in spring.

As shown in Fig. 1, four sampling sites of different types of areas were selected in Baotou. The first sampling site was within the Bayan Obo as a representative MA. Bayan Obo is a typical mining region in north Baotou where a large-scale open-cut (rare earth elements)–Fe–Nb pit is located, and large quantities of raw ore are mined. The second sampling site was a typical smelting area (SA) in west Baotou, which was surrounded by numerous chemical and metallurgical factories, refineries, and power plants. This area is the site to which raw ores from Bayan Obo are transported by railroad for further processing. The third sampling site was a city center area (CA), characterized by a highly dense population, heavy traffic, and commercial activities. Finally, the fourth sampling site was a representative RA of Baotou, located northeast of the CA.

Fig. 1 Map of the study area and sampling sites



Sample collection and analysis

PM₁₀ and PM_{2.5} samples were collected on quartz microfiber filters (MK360, Munktell, Sweden) using a mid-volume aerosol sampler (LaoYing 2030, Qingdao Laoshan Institute of Applied Technology, Qingdao, China) at a flow rate of 100 L min⁻¹. The sampling height was about 1.5 m. At each sampling site, twelve 12-h samples of both PM₁₀ and PM_{2.5} were collected within different time periods between July 25, 2013, and August 30, 2013. Meteorological parameters such as wind speed, wind direction, temperature, and humidity were also recorded at the time of sample collection.

All quartz microfiber filters were dried in a desiccator for 48 h. The filters were weighed before and after aerosol sampling to determine the mass of PM₁₀ and PM_{2.5}. The filters were subsequently sealed in a filter holder and stored at -20 °C until analysis.

To prepare samples for measurement of heavy metals concentrations, each filter was cut into fragments. The fragments were placed in a PTFE crucible and then digested in a mixture of HNO₃, HClO₄ and HF which was heated until about 0.5 mL of colorless solution was obtained. After cooling, the solution was filtered and diluted to a total of 25 mL with Milli-Q[®] water. The concentrations of Cr, Cu, Zn, Pb, and Mn were analyzed by inductively coupled plasma-optical emission spectroscopy (ICP-OES, Optima 5300 DV, Perkin Elmer), and the concentrations of Cd, Co, and Ni were analyzed by inductively coupled plasma-mass spectrometry (ICP-MS, ELAN DRC-e, Perkin Elmer SCIEX). Each measurement was performed in duplicate, and each group had three replicates. National reference samples, replicates, and blanks were also measured to ensure accuracy of the results. The relative error of the results was on average better than 5 %.

Risk assessment

Exposure dose

In this study, the risk assessment model developed by the Environmental Protection Agency (EPA) of the United States was used to evaluate the health risks posed by heavy metals in PM_{2.5}. Considering the variety of physiological characteristics and living styles of Baotou city residents, we divided them into three groups: male (>16 years), female (>16 years) and children (<16 years). Since metal exposure can

occur through direct inhalation, ingestion, and dermal contact, the exposure concentration (EC, µg m⁻³), chemical daily intake (CDI, mg kg⁻¹), and dermal absorbed dose (DAD, mg kg⁻¹) were calculated to assess total exposure dose. EC, CDI, and DAD were calculated according to the Human Health Evaluation Manual (Part A), Supplemental Guidance for Dermal Risk Assessment (Part E), and Supplemental Guidance for Inhalation Risk Assessment (Part F) (EPA 1989, 2004, 2009). The equations are as follows:

$$CDI = \frac{C \times \text{IngR} \times EF \times ED \times CF}{BW \times AT} \quad (1)$$

$$DAD = \frac{C \times SA \times AF \times ABS \times EF \times ED \times CF}{BW \times AT} \quad (2)$$

$$EC = \frac{C \times ET \times EF \times ED}{AT} \quad (3)$$

where *C* stands for the metal concentration in PM_{2.5} (mg kg⁻¹ for CDI and DAD, µg m⁻³ for EC). All of the exposure factors for these models are shown in Table 1.

Risk characterization

The hazard quotient (HQ) was calculated based on exposure dose to assess non-carcinogenic risks posed by airborne metals. The equations are as follows:

$$HQ_{\text{ing}} = CDI/RfDo \quad (4)$$

$$HQ_{\text{inh}} = EC/(RfCi \times 1000 \mu\text{g m}^{-3}) \quad (5)$$

$$HQ_{\text{derm}} = DAD/(RfDo \times GIABS) \quad (6)$$

where RfDo is oral reference dose (mg kg⁻¹ day⁻¹), RfCi is inhalation reference concentrations (µg m⁻³), and GIABS is the gastrointestinal absorption factor. The RfDo, RfCi, and GIABS values for Cr(VI), Ni (refinery dust), Cu, Cd (diet), Pb, Zn (metallic), Co, and Mn (diet) were used from the screening level tables provided by the US EPA (2014).

The hazard index (HI) is equal to the sum of the HQ values for ingestion, inhalation, and dermal contact and represents the total potential non-carcinogenic risks of different pollutants. An HI <1 indicates that there is no significant risk of non-carcinogenic effects, and when HI is >1, a non-carcinogenic effect is likely to exist (EPA 1989).

Table 1 Exposure factors used in assessing health risks

Factor	Definition	Value			Unit	References
		Male	Female	Children		
BW	Average body weight	62.7	54.4	15	kg	Duan (2012)
IngR	Ingestion rate	100	100	200	mg day ⁻¹	EPA (1989)
SA	Surface areas of the skin that contacts the airborne particulates	4220	3820	2160	cm ⁻²	Wang et al. (2008)
AF	Skin adherence factor for the airborne particulates	0.07	0.07	0.2	mg cm ⁻²	EPA (2004)
EF	Exposure frequency	180	180	180	days year ⁻¹	Hu et al. (2012)
ED	Exposure duration	24	24	6	years	EPA (2009)
ET	Exposure time	24			h day ⁻¹	
AT	Averaging time	ED × 365			days	
ABS	Dermal absorption factor	0.001 for Cd, 0.01 for other metals			–	Hu et al. (2012)
CF	Conversion factor	10 ⁻⁶			kg mg ⁻¹	

Cadmium, Cr and Ni (carcinogens) and Pb and Co (probable carcinogens) were chosen for further evaluations as they were classified as problem chemicals by the International Agency for Research on Cancer (IARC 2014). Based on others' previous studies (Greene and Morris 2006; Fang et al. 2013), the individual lifetime lung cancer risk (R_{ic}) was calculated as follows:

$$R_{ic} = (C \times ED \times IUR) / 70 \text{ years} \quad (7)$$

where C is the contaminant concentration ($\mu\text{g m}^{-3}$) and IUR is the inhalation unit risk as defined by the US EPA Integrated Risk Information System (IRIS). The exposure duration (ED) was 92 days year⁻¹ for 70 years in this study. The IUR values for each metal were used from the screening level tables provided by the US EPA (2014).

Due to the fact that heavy metals exposure during childhood may result in a higher lifetime cancer risk than a similar duration exposure during adulthood (EPA 2009), evaluating only R_{ic} may underestimate cancer risks. Thus, it is necessary to take into account age at the time of the exposure. Therefore, age-dependent adjustment factors (ADAFs), as recommended by the US EPA, were used in this study (EPA 2009). Exposure at an age <2 years old requires a tenfold adjustment and at ages 2–16 years old requires a threefold adjustment. No adjustment is needed for exposure at 16 years and older. The equations for lifetime cancer risk were altered to including ADAFs, as follows:

$$\begin{aligned} \text{For a baby (0–2 years): } R_b \\ = R_{ic} \times 10(\text{ADAF}) \times \left(\frac{2 \text{ years}}{70 \text{ years}} \right) \end{aligned} \quad (8)$$

$$\begin{aligned} \text{For a child (2–16 years): } R_c \\ = R_{ic} \times 3(\text{ADAF}) \times \left(\frac{14 \text{ years}}{70 \text{ years}} \right) \end{aligned} \quad (9)$$

$$\begin{aligned} \text{For an adult (> 16 years): } R_a \\ = R_{ic} \times 1(\text{ADAF}) \times \left(\frac{55 \text{ years}}{70 \text{ years}} \right) \end{aligned} \quad (10)$$

The actual lifetime lung cancer risk (R) is the sum of the risk values at each stage in life:

$$R = R_b + R_c + R_a \quad (11)$$

Cancer risks less than 1×10^{-6} are considered negligible by the US EPA.

Results and discussion

The mass concentrations of PM_{10} and $\text{PM}_{2.5}$

The mass concentrations of PM_{10} and $\text{PM}_{2.5}$ determined for the four sampling sites in Baotou city are shown in Fig. 2. The average mass concentrations for $\text{PM}_{2.5}$ and PM_{10} ranged from 26.4 to 129.2 and 71.2 to 318.3 $\mu\text{g m}^{-3}$, respectively, with the highest concentrations found in SA followed by CA, RA, and MA in

Fig. 2 Concentrations of PM_{10} and $PM_{2.5}$ in Baotou ($\mu\text{g m}^{-3}$) (MA mining area, SA smelting area, CA city center area, RA residential area)




Table 2 The average ratio of $PM_{2.5}/PM_{10}$ in different areas

Sites	$PM_{2.5}/PM_{10}$	References
Baotou, China		
SA (smelting area)	0.41	This study
MA (mining area)	0.37	
CA (city center area)	0.51	
RA (residential area)	0.42	
Shenzhen, China	0.73	Lai et al. (2007)
Zhuhai, China	0.71	
Hong Kong, China	0.68	Cheng et al. (2006)
Seoul, Korea	0.73	Kim et al. (2006)
Birmingham, UK	0.66	Yin and Harrison (2008)

that order. Compared to the Air Quality Standard of China ($PM_{2.5} < 75 \mu\text{g m}^{-3}$ and $PM_{10} < 150 \mu\text{g m}^{-3}$), only SA had concentrations of PM_{10} and $PM_{2.5}$ that were much higher than the limit values. In MA, low levels of both PM_{10} and $PM_{2.5}$ were found, which may be attributed to the dry climate and strong wind.

The mass concentration ratio of $PM_{2.5}$ to PM_{10} for MA, SA, CA, and RA were 0.37, 0.41, 0.51, and 0.42, respectively. The average mass concentration ratio was 0.43, meaning that particles between 2.5 and $10 \mu\text{m}$ contribute to weight more than particles below $2.5 \mu\text{m}$. Table 2 summarizes several previously reported $PM_{2.5}/PM_{10}$ values. Compared to other cities, Baotou had a lower $PM_{2.5}/PM_{10}$, indicating a higher coarse particle fraction. Among the different sampling sites, the mass concentration ratios of $PM_{2.5}$ to PM_{10} were ranked in the following order: $MA < SA < RA < CA$. The highest coarse fraction found in MA may be due to the large quantities of dust produced by excavating associated with mining. CA had a relatively high fraction of fine particles compared to the other sites in Baotou. This may be a result of either heavy traffic in the city center or the wind direction

since CA is in the downwind direction of SA, which had the highest level of $PM_{2.5}$.

Heavy metals concentrations in PM_{10} and $PM_{2.5}$

The heavy metal concentrations in $PM_{2.5}$ and PM_{10} of the ambient air are shown in Fig. 3. The highest heavy metal concentration in PM_{10} and $PM_{2.5}$ were found in SA, followed by CA, RA, and MA. It is worth noting that, contrary to expectations, the metal concentrations in particles from MA, a typically active MA, were the lowest. In order to focus on the composition of the PM, the unit of the metal concentration was changed from " $\mu\text{g m}^{-3}$ " to " mg g^{-1} ." This change showed that MA now had the highest concentration of metals in PM_{10} and $PM_{2.5}$ (Fig. 4). The low value when expressed as volume was a result of the low PM concentrations in the air. Although low $PM_{2.5}$ concentrations (weight per volume of air) were measured at MA, comparatively high metal content (mg g^{-1}) of the collected $PM_{2.5}$ was observed, suggests that it is an important source of heavy metals.


Table 3 shows the metal concentration ratios of $PM_{2.5}$ to those in PM_{10} in the four sites of Baotou. According to previous studies (Lee and Hieu 2011; Fang et al. 2013), trace metals are mainly distributed in the fine particles.

Risk assessment

Non-carcinogenic risk assessment

HQ and HI for Cr, Ni, Pb, Cd, Cu, Mn, Co, and Zn in $PM_{2.5}$ samples of each site were calculated using the health risk assessment model of the U.S. EPA. In general, the integrated HI for Baotou residents living in any of the four sampling sites were all higher than the safe level (safe level = 1), indicating a rather high

Fig. 3 Concentrations of eight metals in **a** $PM_{2.5}$ and **b** PM_{10} in Baotou ($\mu g\ m^{-3}$) (MA mining area, SA smelting area, CA city center area, RA residential area)



health risk level from metal exposure in $PM_{2.5}$ (Fig. 5). Residents living in MA and SA faced a higher health risk than residents living in the city center or RA. Among the different groups of residents at the four sites, the integrated HI values increased in the order of male < female < children. The HI values for children were 2–4 times higher than those for adult males and females, indicating that children not only experienced a higher non-carcinogenic risk, but were also more vulnerable to it.

In this study, the HQ values for the three exposure pathways of ingestion, inhalation, and dermal contact varied among the different sites (Fig. 6). There were

notable differences between children and adults. For adults in all sampling sites, the HQ values of the three exposure pathways had the same trends: inhalation > ingestion > dermal contact. The average contribution of HQ_{inh} to HI was 62.6 and 60.2 % for adult males and adult females, respectively. However, for children, the HQ_{ing} was the highest, indicating that ingestion was the most health threatening exposure. Additionally, we found that children faced higher health risks through dermal contact than adults. The HQ_{derm} for adult females and males among all sites were all lower than the safe level (=1), meaning there was not a non-carcinogenic risk posed by heavy metals

Fig. 4 Concentrations of eight metals in **a** $PM_{2.5}$ and **b** PM_{10} in Baotou ($mg\ g^{-1}$) (MA mining area, SA smelting area, CA city center area, RA residential area)




Table 3 The ratios of metal concentrations in $PM_{2.5}$ to that in PM_{10}

	MA (mining area)	SA (smelting area)	CA (city center area)	RA (residential area)
Cr	1.30	1.58	1.23	1.54
Ni	1.72	0.81	0.76	1.17
Cu	1.25	4.51	1.64	2.58
Cd	1.26	3.37	1.73	4.47
Pb	1.38	1.38	2.37	1.10
Zn	1.42	2.66	1.34	2.70
Co	0.85	0.60	0.37	0.88
Mn	0.81	1.57	1.23	2.12




Fig. 5 Non-carcinogenic risks (HI) of residents in Baotou (MA mining area, SA smelting area, CA city center area, RA residential area)

in $PM_{2.5}$. On the contrary, for children, the values of HQ_{derm} were all higher than the safe level and accounted for a larger proportion (20.0 % in average for children, 12.1 % in average for adults) in the integrated HI values.

HI values for Cr, Mn, Cd, and Pb were notably higher than those for other elements (Table 4). In most places, the HI values for Cr and Mn were close to or higher than the safe level, indicating that in Baotou, they might pose non-carcinogenic health risks to both adults and children. Additionally, we found that the Cr HI for children was 2–3 times higher than that for adults at each site, and therefore, Cr might pose higher potential health risks to children. Except for Cr and Mn, the HI values for each selected metal for both adults and children were mostly within the safety range and ranked in the following order: $Pb > Cd > Co > Ni > Cu > Zn$ for CA site, $Cd > Pb > Co > Ni > Cu > Zn$ for all other sites. The higher HI values for Pb in CA site may be attributed to the heavier traffic burden in the city center. Furthermore, the Pb and Cd HI values for children slightly exceeded the safe level, indicating that Pb and Cd pose potential health risks to children and should be studied more extensively.

Lifetime cancer risk assessment

Particulate matter in air, especially in heavily industrialized urban environments, contains a variety of known human carcinogens. In this study, five




Fig. 6 Non-carcinogenic risk distribution of different exposure way for **a** children, **b** adults (female), **c** adults (male) in Baotou (MA mining area, SA smelting area, CA city center area, RA residential area)

carcinogens were investigated. We evaluated the lifetime cancer risks for residents at each site using the mean concentration of each carcinogenic metal in $PM_{2.5}$ and Eq. (11) (Table 5). In all four selected sampling sites in Baotou, the lifetime lung cancer risk was in excess ($>1 \times 10^{-6}$) as posed by the total of five carcinogenic metals (Pb, Cr, Co, Ni and Cd), indicating that carcinogenic risk is not negligible. Among the sites, SA had the highest risks, followed by MA, RA and CA. This indicates that SA residents might face a higher level of cancer risks posed by heavy metals in $PM_{2.5}$.

Among the five selected carcinogenic metals, the cancer risks of Cr, Cd, and Co at all sampling sites were higher than the threshold value 1×10^{-6} . The leading heavy metal was persistently Cr which posed cancer risks 2–3 orders of magnitude higher than the threshold value as well as those posed by other metals. The lifetime cancer risks of Pb and Ni (all sites except for SA) were lower than 1×10^{-6} , implying negligible carcinogenic risk estimates.

Conclusion

The concentration of PM_{10} and $PM_{2.5}$ at all sampling sites except for the SA industrial site was all below the Air Quality Standard of China. SA had the highest metal concentrations per volume of air, while MA had the highest metal concentrations per mass of particles collected. Zn, Mn, Pb, and Cr were the main metal pollutants in both PM_{10} and $PM_{2.5}$. Most selected heavy metals were enriched in the $PM_{2.5}$ fraction at different sampling sites. For the health assessment, all eight selected heavy metals in $PM_{2.5}$ posed non-carcinogenic risks to all groups of residents. Cr, Cd and Co were the most significant contributors to cancer risks in this assessment. Cr had the highest lifetime cancer risk on residents. We hope these results will help raise focus on enforcing more stringent limitations on industrial emissions. Among different groups of people, children experienced the highest health risk in Baotou, followed by adult females and

Table 4 HI values for each non-carcinogenic metal in $PM_{2.5}$ collected in Baotou

	Adults (male)				Adults (female)				Children			
	CA	RA	SA	MA	CA	RA	SA	MA	CA	RA	SA	MA
Cr	0.59	0.62	0.69	1.77	0.63	0.66	0.72	1.91	3.49	3.47	2.96	11.28
Ni	0.03	0.03	0.06	0.04	0.03	0.03	0.06	0.04	0.04	0.05	0.06	0.07
Cu	0.02	0.03	0.03	0.03	0.02	0.04	0.03	0.04	0.14	0.29	0.23	0.27
Cd	0.29	0.49	0.60	0.52	0.30	0.52	0.61	0.56	0.78	1.88	1.29	2.26
Pb	0.30	0.21	0.26	0.31	0.34	0.25	0.30	0.35	2.47	1.78	2.18	2.53
Zn	0.02	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.13	0.11	0.13	0.14
Co	0.05	0.10	0.16	0.16	0.05	0.11	0.17	0.17	0.18	0.48	0.42	0.86
Mn	0.90	1.37	3.26	1.14	0.90	1.38	3.26	1.14	0.96	1.55	3.37	1.35
Total	2.18	2.88	5.06	3.98	2.29	3.00	5.17	4.23	8.20	9.62	10.64	18.77

MA mining area, SA smelting area, CA city center area, RA residential area

Table 5 Lifetime lung cancer risk (*R* values) of residents in Baotou

	MA	SA	CA	RA
Cr	2.24E−04	3.25E−04	1.17E−04	1.56E−04
Ni	7.54E−07	1.13E−06	6.12E−07	6.34E−07
Cd	4.25E−06	7.66E−06	3.35E−06	4.55E−06
Pb	1.25E−07	5.71E−07	3.85E−07	2.61E−07
Co	2.77E−06	5.89E−06	1.46E−06	2.40E−06
Total	2.32E−04	3.40E−04	1.22E−04	1.64E−04

R values higher than safe value (1×10^{-6}) are highlighted in bold

MA mining area, SA smelting area, CA city center area, RA residential area

then adult males. Thus, more attention should be paid on protecting children from pollution hazards.

Acknowledgments This work was sponsored by the National Science Foundation of China (41401591 and 41571473).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Aguilar, J., Dorronsoro, C., Fernández, E., Fernández, J., García, I., Martín, F., et al. (2004). Soil pollution by a pyrite mine spill in Spain: Evolution in time. *Environmental Pollution*, 132(3), 395–401.
- Cancio, J. L., Castellano, A. V., Hernández, M. C., Bethencourt, R. G., & Ortega, E. M. (2008). Metallic species in atmospheric particulate matter in Las Palmas de Gran Canaria. *Journal of Hazardous Materials*, 160(2–3), 521–528.
- Chen, H. W., Chen, W. Y., Chang, C. N., & Chuang, Y. H. (2013). Characterization of particles in the ambience of the high-tech industrial park of central Taiwan. *Aerosol and Air Quality Research*, 13, 699–708.
- Cheng, Y., Ho, K. F., Lee, S. C., & Law, S. W. (2006). Seasonal and diurnal variations of PM_{1.0}, PM_{2.5} and PM₁₀ in the roadside environment of Hong Kong. *China Particuology*, 4(6), 312–315.
- Csavina, J., Field, J., Taylor, M. P., Gao, S., Landázuri, A., et al. (2012). A review on the importance of metals and metalloids in atmospheric dust and aerosol from mining operations. *Science of the Total Environment*, 433, 58–73.
- Duan, X. L. (2012). *Research methods of exposure factors and its application in environmental health risk assessment*. Beijing: Science Press.
- EPA. (1989). Risk assessment guidance for superfund volume I: Human health evaluation manual (Part A). <http://www.epa.gov/oswer/riskassessment/ragsa/>
- EPA. (2004). Risk assessment guidance for superfund volume I: Human health evaluation manual. Part E: Supplemental guidance for dermal risk assessment. <http://www.epa.gov/oswer/riskassessment/rage/index.htm>
- EPA. (2009). Risk assessment guidance for superfund volume I: Human health evaluation manual. Part F: Supplemental guidance for inhalation risk assessment. <http://www.epa.gov/oswer/riskassessment/ragsf/index.htm>
- EPA. (2014). Regional screening level tables. <http://www.epa.gov/region9/superfund/prg/index.html>. Last updated May 2014.
- Espinosa, A. J. F., Rodríguez, M. T., Barragán, F. J., & Sánchez, J. C. J. (2001). Size distribution of metals in urban aerosols in Seville (Spain). *Atmospheric Environment*, 35(14), 2595–2601.
- Fang, W. X., Yang, Y. C., & Xu, Z. M. (2013). PM₁₀ and PM_{2.5} and health risk assessment for heavy metals in a typical factory for cathode ray tube television recycling. *Environmental Science and Technology*, 47, 12469–12476.
- Greene, N. A., & Morris, V. R. (2006). Assessment of public health risks associated with atmospheric exposure to PM_{2.5} in Washington, DC, USA. *International Journal of Environmental Research and Public Health*, 3(1), 86–97.
- Hu, X., Zhang, Y., Ding, Z. H., Wang, T. J., Lian, H. Z., Sun, Y. Y., et al. (2012). Bioaccessibility and health risk of arsenic and heavy metals (Cd, Co, Cr, Cu, Ni, Pb, Zn and Mn) in TSP and PM_{2.5} in Nanjing, China. *Atmospheric Environment*, 57, 146–152.
- IARC. (2014). Agents classified by the IARC monographs (Vol. 1–109). <http://monographs.iarc.fr/ENG/Classification/index.php>
- Kaonga, B. K., & Kgabi, N. A. (2011). Investigation into presence of atmospheric particulate matter in Marikana, mining area in Rustenburg Town, South Africa. *Environmental Monitoring and Assessment*, 178(1), 213–220.
- Kim, K. H., Mishra, V. K., Kang, C. H., Choi, K. C., Kim, Y. J., & Kim, D. S. (2006). The ionic compositions of fine and coarse particle fractions in the two urban areas of Korea. *Journal of Environmental Management*, 78(2), 170–182.
- Klukanová, A., & Rapant, S. (1999). Impact of mining activities upon the environment of the Slovak Republic: Two case studies. *Journal of Geochemical Exploration*, 66(1–2), 299–306.
- Lai, S. C., Zou, S. C., Cao, J. J., Lee, S. C., & Ho, K. F. (2007). Characterizing ionic species in PM_{2.5} and PM₁₀ in four Pearl river delta cities, South China. *Journal of Environmental Sciences*, 19(8), 939–947.
- Lee, B. K., & Hieu, N. T. (2011). Seasonal variation and sources of heavy metals in atmospheric aerosols in a residential

- area of Ulsan, Korea. *Aerosol and Air Quality Research*, 11, 679–688.
- Leili, M., Naddafi, K., Nabizadeh, R., Yunesian, M., & Mesdaghinia, A. (2008). The study of TSP and their heavy metal content in central area of Tehran, Iran. *Air Quality, Atmosphere and Health*, 1(3), 159–166.
- Luís, A. T., Teixeira, P., Almeida, S. F. P., Matos, J. X., & Silva, E. F. (2011). Environmental impact of mining activities in the Lousal area (Portugal): Chemical and diatom characterization of metal-contaminated stream sediments and surface water of Corona stream. *Science of the Total Environment*, 409(20), 4312–4325.
- Pope, C. A., Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Ito, K., et al. (2002). Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *Journal of the American Medical Association*, 287(9), 1132–1141.
- Salomons, W. (1995). Environmental impact of metals derived from mining activities: Processes, predictions, prevention. *Journal of Geochemical Exploration*, 52(1–2), 5–23.
- Sanchez, H. U. R., García, M. D. A., Bejaran, R., Guadalupe, M. E. G., Vázquez, A. W., Toledano, A. C. P., et al. (2009). The spatial–temporal distribution of the atmospheric polluting agents during the period 2000–2005 in the urban area of Guadalajara, Jalisco, Mexico. *Journal of Hazardous Materials*, 165(1–3), 1128–1141.
- Serbula, S. M., Llic, A. A., Kalinovic, J. V., Kalinovic, T. S., & Petrovic, N. B. (2014). Assessment of air pollution originating from copper smelter in Bor (Serbia). *Environmental Earth Sciences*, 71(4), 1651–1661.
- Wang, Z., Liu, S. Q., Chen, X. M., & Lin, C. Y. (2008). Estimates of the exposed dermal surface area of Chinese in view of human health risk assessment. *Journal of Safety and Environment*, 8, 152–156.
- Yin, J. X., & Harrison, R. M. (2008). Pragmatic mass closure study for PM_{1.0}, PM_{2.5} and PM₁₀ at roadside, urban background and rural sites. *Atmospheric Environment*, 42(5), 980–988.
- Zhen, S. (2012). *Air quality evaluation and influence factors analysis of Baotou urban area*. Baotou: School of Economics and Management Inner Mongolia University of Science and Technology.